

September 19, 2023

ERRATA FOR  
**RADIATION DETECTION AND MEASUREMENT:  
CONCEPTS, METHODS, AND DEVICES**

Douglas S. McGregor and J. Kenneth Shultz  
CRC Press, Boca Raton, FL, 2020.  
ISBN13: 978-1439819395  
ISBN10: 1439819394

NOTE: Listed are errors found for the first printing of the first edition.

Location (Discoverer)	As Is	Change to
<b>Chapter 3</b>		
p. 57, Eq. 3.42 (DSM)	$\frac{d^2\psi(x)}{dX^2} - k^2\psi(x) = 0$	$\frac{d^2\psi(x)}{dx^2} - \phi^2\psi(x) = 0$
p. 57, Eq. 3.43 (DSM)	$\frac{d^2\psi(x)}{dX^2} + \phi^2\alpha^2\psi(x) = 0$	$\frac{d^2\psi(x)}{dx^2} + k^2\psi(x) = 0$
p. 57, Eq. 3.44 (DSM)	$\psi_1(x) = Ae^{kx} + Be^{-kx}$	$\psi_1(x) = Ae^{\phi x} + Be^{-\phi x}$
p. 57, Eq. 3.45 (DSM)	$\psi_3(x) = Fe^{kx} + Ge^{-kx}$	$\psi_3(x) = Fe^{\phi x} + Ge^{-\phi x}$
<b>Chapter 6</b>		
p. 225, Example 6.19, line 15 (D. Nichols)	...be $\sigma_{\bar{m}} = \sqrt{7.758}\dots$	...be $\sigma_{\bar{n}} = \sqrt{7.758}\dots$
p. 226, Example 6.20, line 11 (D. Nichols)	$= \left( \frac{g}{t_g} + \frac{b}{t_b} \right)^{1/2} =$	$= \left( \frac{g}{t_G} + \frac{b}{t_B} \right)^{1/2} =$
p. 240, Prob. 7, line 2 (DSM)	$\sigma_n = \frac{1}{N} \sqrt{\sum_{i=1}^N \left( \frac{G_i}{t_{G_i}^2} + \frac{B_i}{t_{B_i}^2} \right)}$	$\sigma_n = \frac{1}{N} \sqrt{\sum_{i=1}^N \left( \frac{G_i}{t_G^2} + \frac{B_i}{t_B^2} \right)}$
<b>Chapter 7</b>		
p. 266, para 3, line 13 (DSM)	...found as $\tau = -n_0/S$ .	...found as $\tau = -S/n_0$ .
p. 267, para 1, line 2 (DSM)	...evaluate $\tau = -n_0/S$ .	...evaluate $\tau = -S/n_0$ .
<b>Chapter 12</b>		
p. 437, Eq. 12.41 (DSM)	$(Ce^{-\gamma b} +$	$(Ce^{-\gamma b} +$
p. 437, Eq. 12.42 (DSM)	$\gamma(Ce^{-\gamma b} +$	$\gamma(Ce^{-\gamma b} +$
p. 454, Ex. 12.2, line 12 (J. Beavers)	$+ \frac{\hbar^2}{3m_t^*}(k_x^2 + k_y^2 + k_z^2 - 2k_xk_y - 2k_xk_z - 2k_yk_z)$	$+ \frac{\hbar^2}{3m_t^*}(k_x^2 + k_y^2 + k_z^2 - k_xk_y - k_xk_z - k_yk_z)$
p. 454, Ex. 12.2, line 12 (J. Beavers)	(all non-diagonal) $- \frac{1}{3} \left( \frac{2}{m_t^*} - \frac{1}{m_l^*} \right)$	(all non-diagonal) $- \frac{1}{3} \left( \frac{1}{m_t^*} - \frac{1}{m_l^*} \right)$
p. 461, para 4, line 9 (J. Beavers)	Here $N_C \equiv 2[2\pi m_e^* kT]/h^2]^{3/2}$ is ...	Here $N_C \equiv 2[[2\pi m_e^* kT]/h^2]^{3/2}$ is ...
p. 470, Ex. 12.3, line 10 (DSM)	Substitution into Eq. (12.141) gives	Substitution into Eq. (12.138) gives
<b>Chapter 13</b>		
p. 488, Eq. 13.9 (DSM)	$\Delta\lambda = \frac{4\pi S\omega c\hbar^2}{E_o^2 - (S\hbar\omega)^2}$	$\Delta\lambda = \frac{4\pi S\omega c\hbar^2}{E_o^2 - (S\hbar\omega)^2}$
p. 490, para 4, line 9 (D. Watson)	the ratio $I(x)/I_o$ is called	the ratio $[I_o - I(x)]/I_o$ is called
p. 494, para 1, line 3 (DSM)	... $d\langle N \rangle/dE = 1/k$ = constant...	... $d\langle N \rangle/dE = k$ = constant...
p. 496, para 3, line 1(DSM)	...electron falls into a luminescent center from its mobile state...	...electron falls from the luminescent center excited state to the ground state...
p. 498, Eq. 13.34 (DSM)	$\% \text{ FWHM} \approx 2\sqrt{2\ln(2)}\sqrt{\frac{1+0.1}{N}} = 2.355\sqrt{\frac{1+0.1}{N}}$	$\% \text{ FWHM} \approx \frac{2\sqrt{2\ln(2)}(100)\sqrt{\frac{1+0.1}{N_p}}}{235.5\sqrt{\frac{1+0.1}{N_p}}} =$
p. 519, para 5, line 3 (DSM)	${}^4\text{He}(\text{n}, {}^3\text{H}){}^4\text{He}$	${}^6\text{Li}(\text{n}, {}^3\text{H}){}^4\text{He}$

(cont.)

Location (Discoverer)	As Is	Change to
<b>Chapter 14</b>		
p. 566, para 1, line 1 (DSM)	...the “Stark effect” to...	...this effect discovered by Austin and Starke to...
p. 571, Eq. 14.6 (DSM)	$\dots \int_0^\infty d^{-\mu r} dr \dots$	$\dots \int_0^\infty e^{-\mu r} dr \dots$
p. 572, para 4, line 4 (DSM)	...near 40 nm (400 angstroms), which...	...near 400 nm (4000 angstroms), which...
p. 573, Eq. 14.11 (DSM)	$T(\lambda) = [(1 - R(\lambda))^2] \dots$	$\tau(\lambda) = T(\lambda) = [(1 - R(\lambda))^2] \dots$
p. 576, Table 14.2, line 8 (for S-24), column 2 (J. Terrell)	SbCs <sub>3</sub>	Na <sub>2</sub> KSb
p. 576, Table 14.2, line 8 (for S-24), column 4 (J. Terrell)	640	620
p. 578, Eq. 14.17 (DSM)	$= \frac{\pi}{3} \left( \frac{2m_e E_F}{\hbar^2} \right)^{3/2}$	$= \frac{8\pi}{3} \left( \frac{2m_e E_F}{\hbar^2} \right)^{3/2}$
p. 590, Eq. 14.37 (K. Huddleston)	$\delta = K \int_0^R \frac{A}{E^n(x)} e^{-\alpha x} dx.$	$\delta = K \int_0^R \frac{A}{E^n(x)} e^{-\alpha x} dx.$
p. 634, Eq. 15.44 (J. Beavers)	$p(x) = p_o + \Delta p \exp \left[ \frac{-x}{\sqrt{D_n \tau_n}} \right]$	$p(x) = p_o + \Delta p \exp \left[ \frac{-x}{\sqrt{D_p \tau_p}} \right]$
p. 644, Ex. 15.2, last line (DSM)	$4.04 \times 10^{-14}$ amperes	$4.04 \times 10^{-10}$ amperes cm <sup>-2</sup>
<b>Chapter 16</b>		
p. 752, Eq. 16.63 (W. McNeil)	$V(y) = \frac{-\rho_c e^{-2y}}{4\kappa\epsilon_0} + C_1 y + C_2.$	$V(y) = \frac{-\rho_c e^{2y}}{4\kappa\epsilon_0} + C_1 y + C_2.$
<b>Chapter 17</b>		
p. 820, Sec.17.2.5, line 7 (D. Nichols)	<sup>113</sup> Cd(n, $\gamma$ ) <sup>113</sup> Cd	<sup>113</sup> Cd(n, $\gamma$ ) <sup>114</sup> Cd
p. 820, Sec.17.2.5, line 11 (D. Nichols)	<sup>113</sup> Cd(n, $\gamma$ ) <sup>113</sup> Cd	<sup>113</sup> Cd(n, $\gamma$ ) <sup>114</sup> Cd
p. 821, para 2, line 1 (D. Nichols)	<sup>113</sup> Cd(n, $\gamma$ ) <sup>113</sup> Cd	<sup>113</sup> Cd(n, $\gamma$ ) <sup>114</sup> Cd
<b>Chapter 19</b>		
p. 970, para 4, line 3 (DSM)	Li <sub>2</sub> B <sub>4</sub> O <sub>7</sub> :Mn has emission ...	Li <sub>2</sub> B <sub>4</sub> O <sub>7</sub> :Cu has emission ...
p. 1008, para 3, line 1 (DSM)	...in 1954 by Gifford...	...in 1964 by Gifford...
p. 1024, Eq. 19.93 (DSM)	$\dots = \frac{2\pi z^2}{137} \left[ \frac{1}{\lambda_2} - \frac{1}{\lambda_1} \right] \sin^2 \theta.$	$\dots = \frac{2\pi \alpha z^2}{137} \left[ \frac{1}{\lambda_2} - \frac{1}{\lambda_1} \right] \sin^2 \theta.$
<b>Chapter 20</b>		
p. 1048, Fig. 20.9 (DSM)	... the product <sup>22</sup> Na transitions ...	... the product <sup>22</sup> Ne transitions ...
p. 1054, Eq. 20.41 (DSM)	$A = \sum_{n=n_1}^{n_2} \left[ C(n) - (n_2 - n_1) \frac{C(n_1) + C(n_2)}{2} \right].$	$A = \sum_{n=n_1}^{n_2} C(n) - \left[ (n_2 - n_1) \frac{C(n_1) + C(n_2)}{2} \right].$
p. 1064, Eq. 20.90 (JKS)	$\frac{dy(x \mathbf{a})}{dB} = f(x, \mu, \tau),$	$\frac{dy(x \mathbf{a})}{dB} = g(x, \mu, \tau),$

(cont.)

Location (Discoverer)	As Is	Change to
p. 1064, Eq. 20.90 (JKS)	$\frac{dy(x \mathbf{a})}{d\mu} = Bf(x, \mu, \tau) \frac{2(x-\mu)}{\tau^2},$	$\frac{dy(x \mathbf{a})}{d\mu} = Bg(x, \mu, \tau) \frac{2(x-\mu)}{\tau^2},$
p. 1064, Eq. 20.90 (JKS)	$\frac{dy(x \mathbf{a})}{d\tau} = f(x, \mu, \tau) \frac{2(x-\mu)^2}{\tau^3},$	$\frac{dy(x \mathbf{a})}{d\tau} = g(x, \mu, \tau) \frac{2(x-\mu)^2}{\tau^3},$
p. 1066, Eq. 20.98 (JKS)	$\frac{dy(x \mathbf{a})}{dB_i} = f(x, \mu_i, \tau_i),$	$\frac{dy(x \mathbf{a})}{dB_i} = g(x, \mu_i, \tau_i),$
p. 1066, Eq. 20.98 (JKS)	$\frac{dy(x \mathbf{a})}{d\mu_i} = Bf(x, \mu_i, \tau_i) \frac{2(x-\mu_i)}{\tau_i^2},$	$\frac{dy(x \mathbf{a})}{d\mu_i} = Bg(x, \mu_i, \tau_i) \frac{2(x-\mu_i)}{\tau_i^2},$
p. 1066, Eq. 20.98 (JKS)	$\frac{dy(x \mathbf{a})}{d\tau_i} = f(x, \mu_i, \tau_i) \frac{2(x-\mu_i)^2}{\tau_i^3},$	$\frac{dy(x \mathbf{a})}{d\tau_i} = g(x, \mu_i, \tau_i) \frac{2(x-\mu_i)^2}{\tau_i^3},$
p. 1066, in solution, line 6 (JKS)	$\mu_1 = 140.$	$\mu_2 = 140.$
p. 1066, in solution, line 10 (JKS)	$\sigma_1 = 2.1298 \pm 0.0157$	$\tau_1 = 2.1298 \pm 0.0157$
p. 1067, in solution, top line (JKS)	$\sigma_2 = 2.8700 \pm 0.0138$	$\tau_2 = 2.8700 \pm 0.0138$

## Chapter 22

p. 1170, Eq 22.61 (DSM)	$\frac{dv_{in}}{dt} - \frac{dv_{out}(t)}{dt} = \frac{1}{C} \frac{dQ(t)}{dt}$	$\frac{dv_{in}(t)}{dt} - \frac{dv_{out}(t)}{dt} = \frac{1}{C} \frac{dQ(t)}{dt}$
p. 1170, line 10 (DSM)	freqencies $\omega \ll 1$ , $G(\omega) \simeq \omega$ so $v_{out} \dots$	freqencies $\omega \ll 1$ and $G(\omega) \simeq \omega\tau$ so $v_{out} \dots$

(cont.)

## Minor Typos

Location (Discoverer)	As Is	Change to
<b>Chapter 1</b>		
p. 5, para 3, line 2 (DSM)	...a gold foil electroscope...	...an electroscope...
p. 5, para 3, line 6 (DSM)	...of the gold foils...	...of the electroscope needle...
p. 5, para 3, line 19 (D. Nichols)	...awarded the 1908 Nobel Prize...	...awarded the 1908 Nobel Prize...
<b>Chapter 3</b>		
p. 59, Eq. 3.57 (DSM)	$\frac{k}{\phi} \left( \left( \frac{kC}{\phi} \sin(ka) - C \cos(ka) \right) = \right)$	$\frac{k}{\phi} \left( \left( \frac{kC}{\phi} \sin(ka) - C \cos(ka) \right) = \right)$
p. 68, Sec. 3.5.4, line 3 (JKS)		The definition of $N_a$ changed in May 2019. See Appendix A.1 for details.
<b>Chapter 4</b>		
p. 144, Eq. 4.154 (DSM)	$\rho R_{\text{CSDA}} \text{ (cm}^2\text{g}^{-1}\text{)} = 10^{a+bx+cx^2}$	$\rho R_{\text{CSDA}} \text{ (in g cm}^{-2}\text{)} = 10^{a+bx+cx^2}$
<b>Chapter 6</b>		
p. 193, Ex. 6.6, line 3 (DSM)	...to ove use...	...to over use...
p. 193, Ex. 6.6, line 4 (DSM)	...possible due...	...possibly due...
p. 216, Eq. 6.74, line 13 (DSM)	$\sigma^2(f) = \sum_{i=1}^N \left( \frac{\partial f(\mu)}{\partial x_i} \right)^2 \sigma_i^2 + \dots$	$\sigma^2(f) = \sum_{i=1}^N \left( \frac{\partial f(\mu)}{\partial x_i} \right)^2 \sigma_i^2 + \dots$
p. 230, Eq. 6.101, line 1 (DSM)	$L_D = L_C + k_\beta \sigma_N =$	$L_D = L_C + k_\beta \sigma_{N_s} =$
p. 230, Eq. 6.102, line 1 (DSM)	$L_D = k_\alpha \sigma_B + k_\beta \sigma_N =$	$L_D = k_\alpha \sigma_0 + k_\beta \sigma_{N_s} =$
p. 230, Para 5, line 3 (DSM)	$(N \ll B)$	$(N_s \ll B)$
p. 230, Eq. 6.103 (DSM)	$L_D = L_C + k_\beta \sigma_N =$	$L_D = L_C + k_\beta \sigma_{N_s} =$
<b>Chapter 9</b>		
p. 320, Ex. 9.1, Line 13 (DSM)	$f N_{O_2} = f \rho_{O_2} N_a / A_{O_2}$	$f_V N_{O_2} = f_V \rho_{O_2} N_a / A_{O_2}$
p. 338, Fig. 9.20 abscissa (DSM)	...arbitray...	...arbitrary...
<b>Chapter 10</b>		
p. 399, Prob. 10.7, line 1 (DSM)	...12.5mm.	...12.5 mm.
<b>Chapter 11</b>		
p. 422, Refs., lines 12,15,18,19 (DSM)	...Elektronenzhlrohr...	...Elektronenzählrohr...
<b>Chapter 12</b>		
p. 437, Eq.12.42 (DSM)	$ig(Ae^{iga} - Be^{-iga})$	$ig(Ae^{iga} - Be^{-iga})$
p. 457, Para 2, line 8 (D. Watson)	...and $E_o$ is...	...and $\mathcal{E}_o$ is ...
p. 462, Eq.12.123 (DSM)	$m_p^{*3/2}$	$m_h^{*3/2}$
p. 471, Ex. 12.3, lines 9, 11, 13 multiple places (J.Beavers)	$\text{cm}^3.$	$\text{cm}^{-3}.$

*(cont.)*

## Minor Typos

Location (Discoverer)	As Is	Change to
p. 472, para 3, line 4 (DSM)	...both side of...	...both sides of...
p. 477, Ex. 12.4, line 1 (J. Beavers)	eV = 1.42 eV	$E_g = 1.42 \text{ eV}$
p. 479, Fig. 12.47 (D. Watson)	point O missing	locate O at origin of $abc$
p. 504, Para 3, line 4 (DSM)	Moszinzki et al. [2003b]	Moszynski et al. [2003b]
p. 504, Para 3, line 11 (DSM)	[Boananami and Rossel, 1952]	[Bonanami and Rossel, 1952]
<b>Chapter 13</b>		
p. 496, para 8, line 4 (DSM)	...times...	...time...
p. 497, Eq. 13.33 (DSM)	$v(Q) \approx \frac{1+\text{var}(G)/\bar{G}^2}{N}$ .	$v(Q) \approx \frac{1+\text{var}(G)/\bar{G}^2}{N\bar{p}} = \frac{1+\text{var}(G)/\bar{G}^2}{N}$ .
p. 498, para 1, line 1 (J. Beavers)	...2002]. Hence, the...	...2002]. Typically $\bar{p} \ll 1$ . Hence, the...
p. 498, Fig. 13.7, line 3 (J. Beavers)	...energy resolution.	...energy resolution for $\bar{p} = 0.25$ .
p. 498, para 2, line 1 (J. Beavers)	This ideal energy resolution is plotted as a solid line in Fig. 13.7, and is compared to measured values of many common inorganic scintillators.	With a common PMT value of $\bar{p} \approx 0.25$ , the ideal energy resolution is plotted and compared to measured values of many inorganic scintillators in Fig. 13.7.
p. 507, para 5, line 2 (DSM)	4.12 g cm <sup>3</sup>	4.12 g cm <sup>-3</sup>
p. 508, para 1, line 2 (DSM)	...with an refractive...	...with a refractive...
p. 508, para 2, line 4 (DSM)	...and the appearance of a luminescent...	...and a luminescent...
p. 525, Table 13.3, line 23, col 10 (DSM)	1,56	1.56
p. 536, para 2, line 13 (DSM)	...about 65°C...	...about 65°...
p. 563, col 2, para 3 (DSM)	SZUPEYCZYNSKI, P.	SZUPRYCZYNSKI, P.
p. 509, para 2, line 8 (DSM)	Is was concluded...	It was concluded...
p. 511, para 1, line 9 (DSM)	[Leroq et al. 2006]	[Lecoq et al. 2006]
p. 518, para 2, line 4 (DSM)	LYSO is 7.4 g cm <sup>-1</sup>	LYSO is 7.12 g cm <sup>-3</sup>
<b>Chapter 14</b>		
p. 575, para 5, line 8 (DSM)	...depends of the thickness...	...depends on the thickness...
p. 579, Fig. 14.12 (D. Watson)	Rh	Rb
p. 581, para 1, line 1 (DSM)	...produce photoexcition.	...produce photoexcitation.
p. 584, para 2, line 10 (DSM)	...forms a NEA boundary...	...forms an NEA boundary...
p. 584, para 2, line 11 (DSM)	...as shown in Fig. 14.16(a).	...as shown in Fig. 14.17(a).

*(cont.)*

## Minor Typos

Location (Discoverer)	As Is	Change to
p. 584, para 2, line 13 (DSM)	...as shown in Fig. 14.16(b).	...as shown in Fig. 14.17(b).
p. 617, para 4, line 5 (DSM)	... $1/M_p$ where $M_p$ ...	... $1/M_h$ where $M_h$ ...
p. 617, Eq. 14.77 (DSM)	...[1 – $\exp(-(\beta - \alpha)x)$ ]...	...[1 – $\exp(-(\beta - \alpha)x)$ ]...
p. 624, Prob. 1, line 1 (DSM)	...Corning 7044 borosilicate window	...Corning 7740 borosilicate window
p. 624, Prob. 3, line 3 (DSM)	$T(\lambda) = \dots$	$\tau(\lambda) = T(\lambda) = \dots$
p. 624, Prob. 6, Line 2 (D. Watson)	...pure CsSb <sub>3</sub> photocathode.	...pure Cs <sub>3</sub> Sb photocathode.
p. 624, Prob. 9, line 2 (DSM)	...from the photodiode?	from the photocathode?
<b>Chapter 15</b>		
p. 639, Para 2, line 5 (D. Watson)	$\dots = kT \ln \left( \frac{n_n p_n}{n_i^2} \right)$	$\dots = kT \ln \left( \frac{n_n p_p}{n_i^2} \right)$
p. 641, third from last line, (J. Beavers)	... $n_{po}$ is the minority hole concentration...	... $n_{po}$ is the minority electron concentration...
p. 644, Ex. 15.2, line 3 (DSM)	mobilities are and...	mobilities are...
p. 645, Fig. 15.11, line 1 (S. Sharma)	...material regiobs...	...material regions...
p. 672, Fig. 15.38, line 2 (DSM)	...yields $n = \dots$	...yields $\check{n} = \dots$
p. 672, para 3, line 1 (DSM)	...of $n, \dots$	...of $\check{n}, \dots$
p. 683, para 5, 2nd to last line (S. Sharma)	...knowledge of $\mu_e \tau_h$ so...	...knowledge of $\mu_e \tau_e^*$ so...
p. 700, problem 2, line 2 (DSM)	$p = 5 \times 10^{15} \text{ cm}^{-3}$ with background dopants $n = 3 \times 10^{15} \text{ cm}^{-3}$	$N_A = 5 \times 10^{15} \text{ cm}^{-3}$ with background dopants $N_D = 3 \times 10^{15} \text{ cm}^{-3}$
p. 701, problem 13, line 1 (DSM)	...values of 50, 25, 0.5, 2.5, 0.5 and 0.05.	...values of 50, 25, 5.0, 2.5, 0.5 and 0.05.
<b>Chapter 16</b>		
p. 726, Ex. 16.1, line 2 (DSM)	...14 to 15 microns...	...16 to 15 microns...
p. 737, Ex. 16.2, line 6 (J. Beavers)	$= \frac{(3.9)(8.854 \times 10^{-14} \text{ F cm}^{-1})}{2 \times 10^{-5} \text{ cm}} =$	$= \frac{(3.9)(8.854 \times 10^{-14} \text{ F cm}^{-1})}{2 \times 10^{-5} \text{ cm}} =$
p. 737, Ex. 16.2, line 8 (DSM)	$Q_{max} = -C_o (V_G - V_T)$	$Q_{max}A = -C_o (V_G - V_T) A$
p. 737, Ex. 16.2, line 8 (DSM)	$= 6.9 \times 10^{-13} \text{ C}$	$= -6.9 \times 10^{-13} \text{ C}$
p. 798, Prob. 10, line 3, (DSM)	electron mobility = $80 \text{ cm}^2 \text{ V}^{-1} \text{ s}^{-1}$ ,	hole mobility = $80 \text{ cm}^2 \text{ V}^{-1} \text{ s}^{-1}$ ,
p. 804, Ref. Hofker (1966), line 2, (DSM)	J.E.J. O-SC BERSKI,	J.E.J. OBERSKI,
p. 805, Ref. C.K. Kim (2009), line 3, (DSM)	2009	1979
p. 809, Ref. Redus, R., (1997), line 1 (DSM)	V. JSC ORDANOV	V. JORDANOV

*(cont.)*

## Minor Typos

Location (Discoverer)	As Is	Change to
<b>Chapter 17</b>		
p. 832, Eq.17.31 (DSM)	... $I_o \exp[-x\sigma_a N_a] = I_0 \exp[-x\Sigma_a]$ .	... $I_0 \exp[-x\sigma_a N_a] = I_0 \exp[-x\Sigma_a]$ .
p. 851, para 2, line 12 (DSM)	... GSO neccessarily have...	... GSO necessarily have...
p. 853, para 3, line 3 (DSM)	...between radition types.	...between radiation types.
p. 856, Fig. 30 (right), in key (DSM)	$^6\text{LiF}$	$^6\text{Li}$
<b>Chapter 18</b>		
p. 943, prob. 4/5 (DSM)	...10-mm-thick <b>5.</b> sample...	...10-mm-thick sample...
<b>Chapter 19</b>		
p. 955, Fig. 19.7 caption (DSM)	<i>all</i> $N_T$	<i>change to</i> $N_t$
p. 955, para 1, line 5 (DSM)	...or $N_{te}(T_0)/N_T$ .	...or $N_{te}(T_0)/N_t$ .
p. 955, para 1, line 6 (DSM)	...of $N_{te}(T_0)/N_T$ .	...of $N_{te}(T_0)/N_t$ .
p. 956, para 1, line 1 (DSM)	...of falling, into a trap...	...of falling into a trap...
p. 957, para 1, line 16 (DSM)	...glow curvs...	...glow curve...
p. 963, para 4, line 6 (DSM)	...generally dosimetry...	...general dosimetry...
p. 965, para 5, line 9 (DSM)	...glow curve ..	...glow curves...
p. 965, para 5, line 10 (DSM)	...ions is significantly different because a glow...	...ions are significantly different because a larger glow...
p. 966, para 3, line 1 (DSM)	...[Kim at al. 2010;...	...[Kim et al. 2010;...
p. 966, para 3, line 4 (DSM)	Kim at al. [2010]...	Kim et al. [2010]...
p. 965, para 6, line 1 (DSM)	...LiF:Mn,Ti...	...LiF:Mg,Ti...
p. 970, para 3, line 1 (DSM)	...of it low...	...of its low...
p. 971, para 1, line 1 (DSM)	... $\times 10^5$ R.	... $10^6$ R.
p. 972, para 7, line 4 (DSM)	...265° that...	...265°C that...
p. 972, para 4, line 5 (DSM)	...incandescensce..	...incandescence...
p. 989, footnote 17, line 3 (DSM)	...and the process...	...and the latter process...
p. 995, para 1, line 2 (DSM)	... $\beta(\delta)$ ...	... $\delta(\beta)$ ...
p. 996, para 2, line 10 (DSM)	...both Eq. (19.43) and Eq. (19.44) vanish...	...both Eq. (19.57) and Eq. (19.58) vanish...

*(cont.)*

## Minor Typos

Location (Discoverer)	As Is	Change to
p. 1002, para 3, line 2 (DSM)	...particles a shown...	...particles as shown...
p. 1003, footnote 22 (DSM)	...Donald Glasser received...	...Donald Glaser received...
p. 1010, para 1, line 1 (DSM)	...so that temperature...	...so that the temperature...
p. 1010, para 2, line 15 (DSM)	...of a radiation interactions...	...of a radiation interaction...
p. 1010, Fig. 19.54, caption (DSM)	...has a absorber...	...has an absorber...
p. 1021, references, col 1, line 3, (DSM)	...Luminescence," <b>81</b> ...	...Luminescence," <i>Rad. Prot. Dos.</i> , <b>81</b> ...
p. 1025, para 3, line 4 (DSM)	...centimete...	...centimeter...
<b>Chapter 20</b>		
p. 1043, line 4 (DSM)	...energy lost from the...	...energy absorbed in the...
p. 1054, para 3, line 3 (DSM)	[Price 2020]	[Press et al. 1992]
p. 1056, para 4, line 4 (DSM)	[Price et al. 1992]	[Press et al. 1992]
p. 1060, para 2, line 2 (DSM)	...(see Appendix C),...	...(see Table 21.4),...
p. 1060, para 2, line 2 (DSM)	...rounded to 520 keV...	...rounded to 352 keV...
p. 1073, Fig. 20.17, abscissa (DSM)	Gamma-Ray Energy	Gamma-Ray Energy (keV)
p. 1076, Ex. 20.5, line 6 (DSM)	$= B_L - (n - n_R) \frac{B_L - B_L}{n_R - n_L} =$	$= B_L - (n - n_R) \frac{B_L - B_R}{n_R - n_L} =$
p. 1092, Eq. 20.149 (DSM)	$C_2 - C_1 = [k_1\sigma(C_2) + k_2\sigma(C_1)]$	$C_2 - C_1 = [k_1\sigma(C_1) + k_2\sigma(C_2)]$
<b>Chapter 22</b>		
p. 1212, para 1, line 4 (DSM)	...detector. while...	...detector, while...
p. 1234, punctuation after eq. 22.226 (DSM)	$P_{max} = \frac{V_{max}^2}{2Z_0}$ .	$P_{max} = \frac{V_{max}^2}{2Z_0},$
p. 1242, Sec A.2, line 7 (JKS)	...is given...	...are given...
<b>Index</b>		
p. 1271, column 2, lines 22, 23, 24, 26, 44, (J. Beavers)	Comptom...	Compton...

*(cont.)*

## Problem Adjustments

To improve the learning experience, these problems are modified to the following:

**Problem 10.2** A coaxial detector is backfilled with P-10 gas to 0.5 atm. The detector has an anode wire with a radius of 25 microns and cathode radius of 1.5 cm. Determine  $r_c$  for an applied voltage of 1500 volts. If the pressure is increased to 2 atm, what is  $r_c$ ?

**Problem 13.7** Suppose you have a NaI:Tl detector with 7% FWHM energy resolution when operated with a 1  $\mu\text{s}$  shaping time at 300 K. What is the expected energy resolution if the temperature is increased to 325 K? Increased to 350 K? Decreased to 250 K?

**Problem 14.5** Determine the thermionic emission current density for a S-11 response PMT at a temperature of 32°C and compare it to that of a S-24 response PMT. Assume that  $A(1 - \alpha_r) \simeq 120 \text{ A cm}^{-2} \text{ K}^{-2}$ .

**Problem 14.6** Determine the electron cutoff energy for the peak wavelength emission from a LaBr<sub>3</sub> detector coupled to a pure CsSb<sub>3</sub> photocathode. Repeat for a Na<sub>2</sub>K<sub>2</sub>Sb photocathode.

**Problem 15.7** You have a Si  $p\nu n$  junction device that is 150 microns wide with  $\nu = 10^{13} \text{ cm}^{-3}$ . Determine the punch through voltage. Determine the breakdown voltage and the punch through breakdown voltage.

**Problem 15.9** Given a 1.2-micron sample of CdS with a shallow trap density of  $10^{15} \text{ cm}^{-3}$ , what is the expected value of  $V_{TFL}$ ? For CdS, the literature value of  $\kappa = \epsilon_s/\epsilon_0 = 8.9$ .


**Problem 18.4/5** Given a thermal neutron beam (0.025 eV) intersecting a 10-mm-thick sample of CLYC:Ce, determine the intrinsic thermal neutron detection efficiency of the CLYC:Ce sample. If the natural Li is replaced with 96% enriched <sup>6</sup>Li, what is the new thermal-neutron detection efficiency?

**Problem 21.5** Potassium and sodium are both Group I elements and thus have similar chemical properties. As a consequence, K is a natural impurity found in NaI:Tl crystals. Estimate the concentration of K in ppm (by mass) to produce two cps in a 4 in  $\times$  5 in (10.16 cm  $\times$  12.7 cm) NaI:Tl detector.


**Problem 21.5** Consider a two-storied house 20  $\times$  20 m in size with walls 8 m high and a basement 3 m deep. The basement floor and walls are of concrete 30-cm thick, and the outside walls are brick 10-cm thick. Plaster 1-cm-thick lines along all walls and the ceilings. Estimate the activity in Bq of <sup>40</sup>K, <sup>226</sup>Ra, and <sup>232</sup>Th in the structural material of the house.

## Corrected Figures

The following figures are described in the erratum above, but for clarity they are reproduced here.




**Figure 9.20**




**corrected Figure 9.20**

*Found by DSM.*




**Figure 12.42**

Found by DSM.




**corrected Figure 12.42**




**Figure 12.45**

**corrected Figure 12.45**

Found by DSM.




**Figure 12.47**




**corrected Figure 12.47**

Found by D. Watson.




**Figure 13.7**




**corrected Figure 13.7**

Found by J. Beavers.




**Figure 13.48**




**corrected Figure 13.48**

Found by DSM.




**Figure 14.12**

Found by D. Watson.




**corrected Figure 14.12**




**Figure 15.38**


Found by DSM.



**corrected Figure 15.38**




**Figure 17.30 (right)**




**corrected Figure 17.30 (right)**

Found by DSM.




**Figure 19.3**




**corrected Figure 19.3**

Found by DSM.



**Figure 20.17**



**corrected Figure 20.17**

Found by DSM.